How to use: Strepsils Lozenges are suitable for adults and children over 6 years. Dissolve one lozenge slowly in the mouth every 2 to 3 hours. Do not take more than 12 lozenges in 24 hours. Do not exceed the stated dose. Not suitable for children under 6 years.

Remember young children can choke on lozenges.

 Like all medicines, this medicine can cause side effects, although not everybody gets them. Side effects may include rash, stomach pain, feeling sick, oral discomfort, rash, allergic reaction.

Consult your doctor if symptoms persist, or anything unusual happens. Do not take for more than 3 days unless told to do so by your doctor.

Contains 0.976g glucose, 1.441g sucrose, 0.3854g fructose per dose. This should be taken into account in patients with diabetes mellitus.

Do not use after the expiry date shown on the pack.

Consult your doctor or pharmacist if you are pregnant or breastfeeding.

KEEP ALL MEDICINES OUT OF THE REACH AND SIGHT OF CHILDREN.

Do not store above 25°C.

PA 979/38/2

Active Ingredients: 2,4-Dichlorobenzyl alcohol 1.2mg, Amylmetacresol 0.6mg.

Also contains: Sucrose, Glucose Syrup, Honey, Tartaric Acid, Peppermint Oil, Terpeneless Lemon Oil and Quinoline Yellow (E104).

Each lozenge contains 1g of glucose and 1.4g of sucrose.

Wording prepared December 2018

PA Holder:

Reckitt Benckiser Reckitt Benckiser

(Ireland) Ltd., Citywest Business Campus,

Dublin 24, Ireland.

Manufacturers: RB Healthcare International Ltd. Nottingham, NG90 2DB, UK and RB NL Brands B.V. Schiphol Blvd 207, 1118 BH Schiphol, NL. Product Queries: Telephone 01-6305429



